

Effect of Tilted Magnetic Field on the Anomalous $H = 0$ Conducting Phase in High-Mobility Si MOSFETs

S. V. Kravchenko, D. Simonian, and M. P. Sarachik

Physics Department, City College of the City University of New York, New York, New York 10031

A. D. Kent

Department of Physics, New York University, New York, New York 10003

V. M. Pudalov

Institute for High Pressure Physics, Troitsk, 142092 Moscow District, Russia

(February 1, 2008)

The suppression by a magnetic field of the anomalous $H = 0$ conducting phase in high-mobility silicon MOSFETs is independent of the angle between the field and the plane of the 2D electron system. In the presence of a parallel field large enough to fully quench the anomalous conducting phase, the behavior is similar to that of disordered GaAs/AlGaAs heterostructures: the system is insulating in zero (perpendicular) field and exhibits reentrant insulator-quantum Hall effect-insulator transitions as a function of perpendicular field. The results demonstrate that the suppression of the low- T phase is related only to the electrons' spin.

PACS numbers: 71.30.+h, 73.40.Qv, 73.40.Hm

According to the one-parameter scaling theory of localization for non-interacting electrons [1], a two-dimensional electron system (2DES) is always insulating at sufficiently large length scales (*i.e.*, in the limit of zero temperature) in the absence of a magnetic field. In high-mobility silicon metal-oxide-semiconductor field-effect transistors (MOSFETs), however, a metal-insulator transition has been observed at a critical electron density, $n_c \sim 10^{11} \text{ cm}^{-2}$, and a $H = 0$ conducting phase has been shown to exist below 1 K [2]. Similar critical behavior has been reported in a p-type SiGe quantum well [3] and in the hole gas in GaAs/AlGaAs heterostructures [4,5]. At low carrier densities, the interaction energy in these systems is more than an order of magnitude larger than the Fermi energy, so that one does not expect the non-interacting theory of localization [1] to be applicable in its simplest form.

In a disordered 2DES, Khmel'nitskii [6] predicted that the extended states that exist at the centers of each Landau level in large perpendicular magnetic fields should "float" up in energy as $H_{\perp} \rightarrow 0$, leading to an insulating phase at $H = 0$. Consistent with this expectation, insulating behavior has been observed in low-density, strongly disordered 2DES in GaAs/AlGaAs heterostructures [7,8]. In contrast, the low-density 2DES in high-mobility Si MOSFETs exhibits quite different behavior. As $H_{\perp} \rightarrow 0$, the extended states shift upward from the centers of the Landau levels [9], as expected. However, instead of "floating" up indefinitely with decreasing magnetic field, the states apparently combine at the Fermi level [9,10], giving rise to anomalous field dependence of ρ_{xx} in small magnetic fields first reported in Ref. [11] and shown in the inset to Fig. 1. This behavior is a puzzle,

and its physical origin has remained unclear.

We have recently shown that the anomalous low-density/low-temperature conducting phase in silicon MOSFETs is suppressed by a magnetic field applied parallel to the 2D plane of the electrons [12,13]: as shown in Fig. 2 in Ref. [12], the resistivity increases by several orders of magnitude as the parallel magnetic field is increased to $H_{\parallel} \sim 20 \text{ kOe}$, above which it saturates to a value that is approximately independent of magnetic field. This prompted us to suggest that the enigmatic behavior in small perpendicular fields is associated with the quenching of a low temperature conducting phase by a perpendicular field (see inset to Fig. 1) just as it is quenched by a parallel field (see Fig. 2 in Ref. [12]). We suggested further that the magnetic field suppression of the anomalous conducting phase in silicon MOSFETs is associated only with the electrons' spin, and is the same for any angle between the field and the 2D plane of the electrons.

From measurements of the resistivity as a function of a magnetic field applied at different angles with respect to the plane of the electrons, we demonstrate in this Letter that: (i) A magnetic field suppresses the anomalous $H = 0$ conducting phase in high-mobility silicon MOSFETs independently of the angle between the field and the plane of the electrons, thereby firmly establishing that the suppression of this phase is associated only with the electrons' spins. (ii) In the presence of a parallel field sufficiently large to quench the anomalous conducting phase in high-mobility silicon samples, the resistivity exhibits as a function of perpendicular field all the now-familiar features found in disordered, low-mobility GaAs/AlGaAs heterostructures [7,8]: a giant negative

magnetoresistance at low H_{\perp} , the quantum Hall effect (QHE) at Landau level filling factors $\nu = 2$ and 1, and insulating behavior at higher H_{\perp} . We also show that: (iii) The suppression of the anomalous conducting phase is not associated with a simple change in mobility or electron density, both of which are essentially unaltered by the magnetic field; and (iv) The multiple valleys that are




FIG. 1. Resistivity as a function of the total magnetic field for a high-mobility sample B, at $T = 0.36$ K and $n_s = 1.0 \times 10^{11} \text{ cm}^{-2}$, for nine angles ϕ between the magnetic field and the inversion layer. ρ_{xx} deviates from the “main” curve at smaller magnetic fields as ϕ is increased. (The inset shows the resistance of sample A at $T = 35$ mK as a function of a perpendicular magnetic field. At the electron density of $n_s = 9.3 \times 10^{10} \text{ cm}^{-2}$ this sample is in the conducting state at $H = 0$. QHE resistivity minima at filling factors $\nu = 1, 2$, and 6 are shown by arrows. The resistivity decreases rapidly as $H \rightarrow 0$, approaching a finite value contrary to the expected insulating behavior).

peculiar to the conduction band of silicon are not responsible for the low-temperature conducting phase, which is suppressed the same way by a field applied at any angle.

The three silicon MOSFET samples used for these studies have peak mobilities at 4.2 K of $\mu_{4.2\text{K}}^{\text{max}} \approx 30,000 \text{ cm}^2/\text{Vs}$ (sample A), $25,000 \text{ cm}^2/\text{Vs}$ (sample B), and $8,000 \text{ cm}^2/\text{Vs}$ (sample C). Four-terminal DC transport measurements were taken as a function of a magnetic field applied at different angles with respect to the

plane of the electrons. Two Si MOSFET samples were measured in a pumped ^3He system equipped with a 12-Tesla magnet and a manual sample rotator. Sample A was studied in a dilution refrigerator in a magnetic field oriented perpendicular to the 2D plane. Excitation currents were between 0.01 nA and 10 nA; care was taken to ensure measurements were in the linear $I - V$ regime.

For a gate voltage that placed sample B in the conducting state at $H = 0$ with a resistivity of $\approx 10 \text{ k}\Omega$ at 360 mK, Fig. 1 shows the diagonal resistivity, ρ_{xx} , as a function of a magnetic field applied at different angles with respect to the plane of the 2DES. For all angles, $\rho_{xx}(H)$ follows approximately the same curve up to some value of magnetic field, above which orbital effects leading to QH oscillations become dominant. The resistivity deviates from the “main” curve at smaller magnetic fields as the angle between the field and the plane is increased: the larger perpendicular component causes stronger orbital effects which become dominant at a lower total field. We note that small differences in $\rho_{xx}(H)$ at $H \sim 10 \text{ kOe}$ are associated with the emergence of a QHE minimum at filling factor $\nu = 6$ [10], which deepens as the perpendicular component of the field gets larger. The important feature is that the magnetoresistance is the same at all angles up to some field above which it is overwhelmed by orbital effects. The anomalous $H = 0$ conducting phase is thus suppressed in the same manner by a magnetic field applied at any angle.

This provides evidence that the conduction band valleys in silicon do not play an important role. It has been shown [14] that a field applied parallel to the plane of the 2DES in silicon MOSFETs does not affect the splitting of two conduction band valleys. This splitting is enhanced in a perpendicular field due to exchange interactions, and is therefore expected to be a function of field orientation. The absence of any angular dependence implies that valley-splitting is not responsible for the suppression of the low-temperature conducting phase by a magnetic field. We thus arrive at the important conclusion that it is the electrons’ spin that plays a crucial role. Indeed, among the theoretical suggestions that have been offered as possible explanations of the conducting phase at $H = 0$ [15,16,17,18,19,20,21], several involve electron spins [15,17,18,19].

We now verify explicitly that a magnetic field does not drive the sample into the insulating phase by simply reducing the electron mobility [22], or by reducing the electron density below its critical value. Fig. 2 shows $\mu_{4.2\text{K}}$ of high-mobility sample B as a function of electron density in $H = 0$ and in the presence of a parallel magnetic field, $H_{\parallel} = 30 \text{ kOe}$. These data establish that the mobility is essentially unaltered by a magnetic field.

The inset to Fig. 2 shows the resistance as a function of the perpendicular component of the magnetic field,




FIG. 2. Mobility *vs* electron density for sample B at $T = 4.2$ K in zero magnetic field (open symbols) and $H_{||} = 30$ kOe (closed symbols). The inset shows R_{xx} as a function of H_{\perp} for four angles between the field and 2D plane; $T = 0.36$ K and $n_s = 1.0 \times 10^{11} \text{ cm}^{-2}$.

$H_{\perp} = H \sin \phi$, as the total field H is swept at four different fixed angles with respect to the electron plane. Note that the parallel field, $H_{||} = H \cos \phi$, varies along each curve and is different for different angles ϕ . The QHE minima occur at the same H_{\perp} for all angles, corresponding to different values of the total field. This observation establishes that the magnetic field does not change the electron density in the inversion layer. The dramatic growth with angle of the ρ_{xx} maximum at $H_{\perp} \sim 15$ kOe can be understood by noting that the $H = 0$ conducting state is quenched independently of the field orientation: at a fixed $H_{\perp} \approx 15$ kOe, the total field increases with decreasing ϕ , $H = H_{\perp}(\sin \phi)^{-1}$, driving the sample closer to the insulating state. Note that an anomalous growth with $H_{||}$ of the resistance peak between $\nu = 1$ and $\nu = 3$ has been observed in a p-Si/SiGe heterostructure [24] and was attributed by the authors to the dependence of the “insulating state width on the ratio between spin and cyclotron splittings”. We remark that the observation of a $H = 0$ conducting state similar to that in high-mobility Si MOSFETs in this system [3] suggests that the strong enhancement of the resistivity in p-Si/SiGe [24] may instead be due to the magnetic field suppression of the anomalous conducting state in the same way as in Si MOSFETs.




FIG. 3. ρ_{xx} of sample B as a function of H_{\perp} for different values of the parallel magnetic field; $T = 0.36$ K and $n_s = 1.0 \times 10^{11} \text{ cm}^{-2}$. The inset shows $\rho_{xx}(H_{\perp})$ for a low-mobility sample C; $T = 0.36$ K and $n_s = 2.1 \times 10^{11} \text{ cm}^{-2}$.

The diagonal resistivity, ρ_{xx} , is plotted as a function of H_{\perp} in several fixed parallel magnetic fields in Fig. 3. The lowest curve corresponds to $H_{||} = 0$ and exhibits the anomalous behavior of high-mobility Si MOSFETs [11]. The peak at $H_{\perp} \approx 15$ kOe is considerably smaller than that shown in the inset to Fig. 1 because of the higher measuring temperature (360 mK *vs* 35 mK). The highest curve is the magnetoresistance of the sample in the insulating state, obtained by quenching the $H = 0$ conducting state with a parallel field of 34 kOe. In the “quenched” phase, high-mobility Si MOSFETs display the familiar reentrant behavior found in disordered, weakly interacting GaAs/AlGaAs heterostructures (see, *e.g.*, Fig. 2 in Ref. [7]): the system has an initial large negative magnetoresistance, exhibits the quantum Hall effect at $\nu = 2$ and 1, and becomes again insulating at $H \gtrsim 42$ kOe. (However, the initial decrease in resistivity is considerably less sharp than in disordered GaAs/AlGaAs.) The gradual disappearance of the anomalous conducting phase is apparent at intermediate fields. Note that above $H_{\perp} \sim 20$ kOe all the data collapse onto a single curve. This confirms once again that the anomalous phase is quenched by a magnetic field applied in any direction (including perpendicular).

The inset to Fig. 3 shows the diagonal magnetoresistivity ρ_{xx} of the relatively low-mobility sample C in a perpendicular field. No $H = 0$ conducting phase was found in this sample. It is strongly insulating at $H_{\perp} = 0$, and there is an appreciable negative magnetoresistance for $H_{\perp} \lesssim 30$ kOe. The $\nu = 1$ and $\nu = 2$ QHE minima in ρ_{xx} are evident, followed at higher field by a transition to an insulator due to the crossing of the last extended state through the Fermi level at $H_{\perp} \gtrsim 65$ kOe. It is interesting that ρ_{xx} vs H_{\perp} for sample C is qualitatively similar to the behavior of the high-mobility sample B in a partially quenched phase. Moreover, sample C exhibits a strong positive magnetoresistance as a function of H_{\parallel} (not shown). This suggests that the anomalous low-temperature phase that is so evident in high-mobility samples is also present in a modified, partially quenched form in low-mobility, disordered samples.

In conclusion, we have shown that the suppression by a magnetic field of the $H = 0$ conducting phase in high-mobility Si MOSFETs does not depend on the angle the field makes with the 2D electron plane. This provides strong evidence that valley splitting does not play an important role, and that the quenching of the anomalous conducting phase in two dimensions is associated with the electrons' spin. We have also demonstrated explicitly that the suppression of the conductivity is not associated with a simple change in sample mobility or electron density, both of which are essentially unaffected by magnetic field. In the presence of large parallel field, the "quenched" phase in high-mobility silicon MOSFETs exhibits the reentrant behavior of disordered, weakly interacting GaAs/AlGaAs heterostructures: a large negative magnetoresistance and reentrant insulator-QHE-insulator transitions [7,8].

We are indebted to Robert Wheeler for generously supplying MOSFET devices fabricated in his laboratory. We thank Veronika Simonian for help in analyzing data using MathematicaTM, Mark Ofitserov for assistance with the experimental equipment, and Dan Shahar for useful discussions. This work was supported by the US Department of Energy under Grant No. DE-FG02-84ER45153. A. K. was supported by NYU and an NYU Research Challenge Fund Grant. V. P. was supported by RFBR (grant 97-02-17387) and by INTAS.

[1] E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).
[2] S. V. Kravchenko, G. V. Kravchenko, J. E. Furneaux, V. M. Pudalov, and M. D'Iorio, Phys. Rev. B **50**, 8039 (1994);

S. V. Kravchenko, W. E. Mason, G. E. Bowker, J. E. Furneaux, V. M. Pudalov, and M. D'Iorio, Phys. Rev. B **51**, 7038 (1995); S. V. Kravchenko, D. Simonian, M. P. Sarachik, W. Mason, and J. E. Furneaux, Phys. Rev. Lett. **77**, 4938 (1996).
[3] P. T. Coleridge, R. L. Williams, Y. Feng, and P. Zawadzki, to be published in Phys. Rev. B, Rapid Communications; see also preprint cond-mat/9708118.
[4] Y. Hanein, U. Meirav, D. Shahar, C. C. Li, D. C. Tsui, and H. Shtrikman, cond-mat/9709184.
[5] M. Y. Simmons, A. R. Hamilton, M. Pepper, E. H. Linfield, P. D. Rose, and D. A. Ritchie, cond-mat/9709240.
[6] D. E. Khmel'nitskii, Pis'ma Zh. Eksp. Teor. Fiz. **38**, 454 (1983) [JETP Lett. **38**, 552 (1983)]; Phys. Lett. A **106**, 182 (1984); see also R. B. Laughlin, Phys. Rev. Lett. **52**, 2304 (1984); S. Kivelson, D.-H. Lee, and S. C. Zhang, Phys. Rev. B **46**, 2223 (1992).
[7] H. W. Jiang, C. E. Johnson, K. L. Wang, and S. T. Hannahs, Phys. Rev. Lett. **71**, 1439 (1993).
[8] T. Wang *et al*, Phys. Rev. Lett. **72**, 709 (1994); R. J. F. Hughes *et al*, J. Phys. Condens. Matter **6**, 4763 (1994); D. Shahar, D. C. Tsui, and J. E. Cunningham, Phys. Rev. B **52**, R14 372 (1995); I. Glozman, C. E. Johnson, and H. W. Jiang, Phys. Rev. Lett. **74**, 594 (1995).
[9] A. A. Shashkin, G. V. Kravchenko, and V. T. Dolgoplov, Pis'ma Zh. Eksp. Teor. Fiz. **58**, 215 (1993) [JETP Lett. **58**, 220 (1993)].
[10] V. M. Pudalov, M. D'Iorio, and J. W. Campbell, Surf. Sci. **305**, 107 (1994).
[11] M. D'Iorio, V. M. Pudalov, and S. G. Semenchinsky, Phys. Lett. A **150**, 422 (1990).
[12] D. Simonian, S. V. Kravchenko, M. P. Sarachik, and V. M. Pudalov, Phys. Rev. Lett. **79**, 2304 (1997).
[13] V. M. Pudalov, G. Brunthaler, A. Prinz, and G. Bauer, Pis'ma Zh. Eksp. Teor. Fiz. **65**, 887 (1997) [JETP Lett. **65**, 932 (1997)].
[14] R. J. Nicholas, K. von Klitzing, and Th. Englert, Solid State Commun. **34**, 51 (1980), see also T. Ando, A. B. Fowler, and F. Stern, Rev. Mod. Phys. **54**, 437 (1982) and ref's therein.
[15] A. M. Finkel'shtein, Zh. Eksp. Teor. Fiz. **84**, 168 (1983) [Sov. Phys. JETP **57**, 97 (1983)].
[16] V. Dobrosavljević, E. Abrahams, E. Miranda, and S. Chakravarty, Phys. Rev. Lett. **79**, 455 (1997).
[17] D. Belitz and T. R. Kirkpatrick, cond-mat/9705023.
[18] P. Phillips, Y. Wan, I. Martin, S. Knysh, and D. Dalidovich, submitted to Nature (London); cond-mat/9709168.
[19] V. M. Pudalov, cond-mat/9707076.
[20] C. Bulutay and M. Tomak, cond-mat/9707339.
[21] F.-C. Zhang and T. M. Rice, cond-mat/9708050.
[22] Disappearance of the metallic phase with decreasing mobility has been reported in the first paper in Ref. [2] and demonstrated on a single sample with variable mobility in Ref. [23].
[23] D. Popović, A. B. Fowler, and S. Washburn, Phys. Rev. Lett. **79**, 1543 (1997).
[24] S. I. Dorozhkin, C. J. Emeleus, T. E. Whall, and G. Landwehr, Phys. Rev. B **52**, R11 638 (1995).